

HPV
also known as Human Papillomavirus

As parents, you do everything you can to protect your children's health for now and for the future. Today, there is a strong weapon to prevent several types of cancer in our kids: the HPV vaccine.

HPV and Cancer

HPV is short for Human Papillomavirus, a common virus. In the United States each year, there are about 17,500 women and 9,300 men affected by HPV-related cancers. Many of these cancers **could be prevented with vaccination.** In both women and men, HPV can cause anal cancer and mouth/throat (oropharyngeal) cancer. It can also cause cancers of the cervix, vulva and vagina in women; and cancer of the penis in men.

For women, screening is available to detect most cases of cervical cancer with a Pap smear. Unfortunately, there is no routine screening for other HPV-related cancers for women or men, and these cancers can cause pain, suffering, or even death. **That is why a vaccine that prevents most of these types of cancers is so important.**

More about HPV

HPV is a virus passed from one person to another during skin-to-skin sexual contact, including vaginal, oral, and anal sex. HPV is most common in people in their late teens and early 20s. Almost all sexually active people will get HPV at some time in their lives, though most will never even know it.

Most of the time, the body naturally fights off HPV, before HPV causes any health problems. But in some cases, the body does not fight off HPV, and HPV can cause health problems, like cancer and genital warts. Genital warts are not a life-threatening disease, but they can cause emotional stress, and their treatment can be very uncomfortable. About 1 in 100 sexually active adults in the United States have genital warts at any given time.

HPV vaccination is recommended for preteen girls and boys at age 11 or 12 years

All preteens need HPV vaccination so they can be protected from HPV infections that cause cancer. Teens and young adults who didn't start or finish the HPV vaccine series also need HPV vaccination. Young women can get HPV vaccine until they are 27 years old and young men can get HPV vaccine until they are 22 years old. Young men who have sex with other men or who have weakened immune systems can also get HPV vaccine until they are 27.

HPV vaccination is a series of shots given over several months. The best way to remember to get your child all of the shots they need is to make an appointment for the remaining shots before you leave the doctor's office or clinic.

Is the HPV vaccine safe?

Yes. HPV vaccination has been studied very carefully and continues to be monitored by CDC and the Food and Drug Administration (FDA). No serious safety concerns have been linked to HPV vaccination. **These studies continue to show that HPV vaccines are safe.**

The most common side effects reported after HPV vaccination are mild. They include pain and redness in the area of the arm where the shot was given, fever, dizziness, and nausea. Some preteens and teens may faint after getting a shot or any other medical procedure. Sitting or lying down for about 15 minutes after getting shots can help prevent injuries that could happen if your child were to fall while fainting. ▶



Why does my child need this now?

HPV vaccines offer the best protection to girls and boys who complete the series and have time to develop an immune response **before** they begin sexual activity with another person. This is not to say that your preteen is ready to have sex. In fact, it's just the opposite—it's important to get your child protected before you or your child have to think about this issue. The immune response to this vaccine is better in preteens, and this could mean better protection for your child. ❖

DISTRIBUTED BY:



Serious side effects from HPV vaccination are rare. Children with severe allergies to yeast or latex shouldn't get certain HPV vaccines. Be sure to tell the doctor or nurse if your child has any severe allergies.

Help paying for vaccines

The Vaccines for Children (VFC) program provides vaccines for children ages 18 years and younger who are uninsured, Medicaid-eligible, or American Indian/Alaska Native. Learn more about the VFC program at

www.cdc.gov/Features/VFCprogram/

Whether you have insurance, or your child is VFC-eligible, some doctors' offices may also charge a fee to give the vaccines. ■

Jacquelyn's story: "I was healthy—and got cervical cancer."

When I was in my late 20's and early 30's, in the years before my daughter was born, I had some abnormal Pap smears and had to have further testing. I was told I had the kind of HPV that can cause cancer and mild dysplasia.

For three more years, I had normal tests. But when I got my first Pap test after my son was born, they told me I needed a biopsy. The results came back as cancer, and my doctor sent me to an oncologist. Fortunately, the cancer was at an early stage. My lymph nodes were clear, and I didn't need radiation. But I did need to have a total hysterectomy.

My husband and I have been together for 15 years, and we were planning to have more children. We are so grateful for our two wonderful children, but we were hoping for more—which is not going to happen now.

The bottom line is they caught the cancer early, but the complications continue to impact my life and my family. For the next few years, I have to get pelvic exams and Pap smears every few months, the doctors measure tumor markers, and I have to have regular x-rays and ultrasounds, just in case. I have so many medical appointments that are taking time away from my family, my friends, and my job.

Worse, every time the phone rings, and I know it's my oncologist calling, I hold my breath until I get the results. I'm hopeful I can live a full and healthy life, but cancer is always in the back of my mind.

In a short period of time, I went from being healthy and planning more children to all of a sudden having a radical hysterectomy and trying to make sure I don't have cancer again. It's kind of overwhelming. And I am one of the lucky ones!

Ultimately I need to make sure I'm healthy and there for my children. I want to be around to see their children grow up.

I will do everything to keep my son and daughter from going through this. I will get them both the HPV vaccine as soon as they turn 11. I tell everyone—my friends, my family—to get their children the HPV vaccine series to protect them from this kind of cancer. ❖



What about boys?

HPV vaccine is for boys too! This vaccine can help prevent boys from getting infected with the types of HPV that can cause cancers of the mouth/throat, penis and anus. The vaccine can also help prevent genital warts. HPV vaccination of males is also likely to benefit females by reducing the spread of HPV viruses.

Learn more about HPV and HPV vaccine at www.cdc.gov/hpv

For more information about the vaccines recommended for preteens and teens:

800-CDC-INFO (800-232-4636)
www.cdc.gov/vaccines/teens

HPV (Human Papillomavirus) Vaccine: What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de Información Sobre Vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

HPV vaccine prevents infection with human papillomavirus (HPV) types that are associated with many cancers, including:

- **cervical cancer** in females,
- **vaginal and vulvar cancers** in females,
- **anal cancer** in females and males,
- **throat cancer** in females and males, and
- **penile cancer** in males.

In addition, HPV vaccine prevents infection with HPV types that cause **genital warts** in both females and males.

In the U.S., about 12,000 women get cervical cancer every year, and about 4,000 women die from it. HPV vaccine can prevent most of these cases of cervical cancer.

Vaccination is not a substitute for cervical cancer screening. This vaccine does not protect against all HPV types that can cause cervical cancer. Women should still get regular Pap tests.

HPV infection usually comes from sexual contact, and most people will become infected at some point in their life. About 14 million Americans, including teens, get infected every year. Most infections will go away on their own and not cause serious problems. But thousands of women and men get cancer and other diseases from HPV.

2 HPV vaccine

HPV vaccine is approved by FDA and is recommended by CDC for both males and females. It is routinely given at 11 or 12 years of age, but it may be given beginning at age 9 years through age 26 years.

Most adolescents 9 through 14 years of age should get HPV vaccine as a two-dose series with the doses separated by 6-12 months. People who start HPV vaccination at 15 years of age and older should get the vaccine as a three-dose series with the second dose given 1-2 months after the first dose and the third dose given 6 months after the first dose. There are several exceptions to these age recommendations. Your health care provider can give you more information.

3 Some people should not get this vaccine

- Anyone who has had a severe (life-threatening) allergic reaction to a dose of HPV vaccine should not get another dose.
- Anyone who has a severe (life threatening) allergy to any component of HPV vaccine should not get the vaccine.

Tell your doctor if you have any severe allergies that you know of, including a severe allergy to yeast.

- HPV vaccine is not recommended for pregnant women. If you learn that you were pregnant when you were vaccinated, there is no reason to expect any problems for you or your baby. Any woman who learns she was pregnant when she got HPV vaccine is encouraged to contact the manufacturer's registry for HPV vaccination during pregnancy at 1-800-986-8999. Women who are breastfeeding may be vaccinated.
- If you have a mild illness, such as a cold, you can probably get the vaccine today. If you are moderately or severely ill, you should probably wait until you recover. Your doctor can advise you.

4 Risks of a vaccine reaction

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own, but serious reactions are also possible.

Most people who get HPV vaccine do not have any serious problems with it.

Mild or moderate problems following HPV vaccine:

- Reactions in the arm where the shot was given:
 - Soreness (about 9 people in 10)
 - Redness or swelling (about 1 person in 3)
- Fever:
 - Mild (100°F) (about 1 person in 10)
 - Moderate (102°F) (about 1 person in 65)
- Other problems:
 - Headache (about 1 person in 3)



Problems that could happen after any injected vaccine:

- People sometimes faint after a medical procedure, including vaccination. Sitting or lying down for about 15 minutes can help prevent fainting, and injuries caused by a fall. Tell your doctor if you feel dizzy, or have vision changes or ringing in the ears.
- Some people get severe pain in the shoulder and have difficulty moving the arm where a shot was given. This happens very rarely.
- Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/.

5 What if there is a serious reaction?

What should I look for?

Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a **severe allergic reaction** can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

If you think it is a **severe allergic reaction** or other emergency that can't wait, call 9-1-1 or get to the nearest hospital. Otherwise, call your doctor.

Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your health care provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/hpv

Vaccine Information Statement HPV Vaccine

12/02/2016

42 U.S.C. § 300aa-26

Office Use Only



HPV Vaccine is Safe – (Gardasil)

04/08/2016 CS256663A

What are HPV Vaccines?

HPV vaccines protect against certain cancers caused by human papillomavirus (HPV) infection. HPV infection can cause cervical, vaginal, and vulvar cancers in women and penile cancer in men. HPV can also cause anal cancer, throat (oropharyngeal) cancer, and genital warts in both men and women. There are currently three HPV vaccines available for use in the United States. This fact sheet summarizes what we know about the safety of Gardasil, one of the available HPV vaccines.

How Do I Know HPV Vaccine is Safe?

As with all approved vaccines, CDC and the Food and Drug Administration (FDA) closely monitor the safety of HPV vaccine to identify **adverse events** and **side effects**. Pre-licensure clinical trials and data collected after the vaccine was made available show that it is very safe.

Adverse event: a health problem that happens after vaccination that may or may not be caused by a vaccine.

Side effect: a health problem that has been shown to be linked to a vaccine by scientific studies.

What Are the Side Effects?

HPV vaccine is very safe, and it is effective at protecting against some HPV types that cause cancer. Vaccines, like any medicine, can have side effects. Many people who get HPV vaccine have no side effects at all. Some people report having very mild side effects, like a sore arm from the shot. The most common side effects are usually mild.

Common Side Effects of HPV Vaccines

- Pain, redness, or swelling in the arm where the shot was given
- Headache or feeling tired
- Fever
- Nausea
- Muscle or joint pain

Understanding HPV Vaccine Safety Studies and Monitoring

It is important to understand the following when reading about HPV vaccine safety studies:

Anyone can report side effects and adverse events.

CDC and FDA maintain a vaccine safety monitoring system called the [Vaccine Adverse Event Reporting System \(VAERS\)](#). VAERS accepts reports from anyone, including doctors, patients, and parents. While VAERS provides useful information on vaccine safety, the data have limitations. It is generally not possible to use VAERS to determine whether a vaccine caused an adverse event.

HPV vaccine has many of the same, mild side effects as other vaccines.

Common, mild side effects reported during HPV vaccine safety studies include pain in the arm where the shot was given, fever, dizziness and nausea. These are similar to side effects seen with other vaccines.

Some preteens and teens might faint after getting the HPV vaccine or any shot. People should sit or lie down for about 15 minutes after getting a shot. This can help prevent fainting.

CDC has carefully studied the risks of HPV vaccination.

HPV vaccination is recommended because the benefits, such as prevention of cancer, far outweigh the risks of possible side effects.

Benefits	Potential Risks
Cancer Prevention	Chance of fainting
Cervical, vaginal, and vulvar cancer in women	Pain, redness, or swelling in the arm where the shot was given
Anal cancer in men and women	
Likely penile cancer in men	
Likely oropharyngeal cancer in women and men	



AMERICAN ACADEMY OF
FAMILY PHYSICIANS
STRONG MEDICINE FOR AMERICA

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

What Do the Studies Say?

Scientists at CDC and FDA continuously monitor the safety of HPV vaccine. While monitoring activities help identify possible side effects and adverse events, they do not prove the side effects were caused by Gardasil.

- [Monitoring](#) by CDC and FDA in 2009 revealed most side effects reported after receiving HPV vaccine were non-serious, including: fainting; dizziness; nausea; headache; and pain, swelling, or redness in the arm where the shot was given.

Formal studies have also looked at whether or not specific adverse events can be linked to Gardasil:

- A 2011 [study](#) found women and girls who received Gardasil were no more at risk of allergic reactions, anaphylaxis (severe allergic reaction), Guillain-Barré Syndrome (GBS), stroke, blood clots, appendicitis, or seizures than those who were unvaccinated or who received other vaccines.
- A 2012 [study](#) that looked at when adverse events occur found Gardasil may be associated with skin infections where the shot is given during the two weeks after vaccination and fainting on the day the shot is received.
- A 2013 [study](#) that included almost 1 million girls found Gardasil was not associated with blood clots or adverse events related to the autoimmune and brain systems.
- A 2014 [study](#) that included over 1 million women found Gardasil was not associated with venous thromboembolism, also called VTE or blood clots.

Several studies have shown that there is no relationship between Gardasil and [autoimmune disorders](#):

- A 2012 [study](#) and a 2014 [study](#) both found women and girls who received the Gardasil shot were not more likely to develop autoimmune disorders than those who were unvaccinated.
- A 2015 [study](#) found women and girls who received Gardasil were not more likely than those who were unvaccinated to develop multiple sclerosis (MS) or other similar diseases.

Gardasil is not recommended during pregnancy. However, some women may receive the Gardasil shot before realizing they are pregnant. There have been several studies that found pregnant women who received Gardasil did not experience any problems:

- A 2015 [study](#) found no safety concerns for pregnant women who received Gardasil, or for their babies.
- The Gardasil Pregnancy Registry, maintained by the manufacturer, received many reports of pregnant women who were vaccinated, and found no evidence that the vaccine affects fertility, pregnancy, or the health of the baby.

Related Scientific Articles

Arnheim-Dahlström L, Pasternak B, Svanström H, Sparén P, Hviid A. [Autoimmune, neurological, and venous thromboembolic adverse events after immunisation of adolescent girls with quadrivalent human papillomavirus vaccine in Denmark and Sweden: Cohort study](#). *BMJ*. 2013 Oct;347:f5906.

Chao C, Klein NP, Velicer CM, Sy LS, Slezak JM, Takhar H, et al. [Surveillance of autoimmune conditions following routine use of quadrivalent human papillomavirus vaccine](#). *J Intern Med*. 2012 Feb; 271(2):193-203. Epub 2011 Nov.

Gee J, Naleway A, Shui I, Baggs J, Yin R, Li R, et al. [Monitoring the safety of quadrivalent human papillomavirus vaccine: Findings from the Vaccine Safety Datalink](#). *Vaccine*. 2011 Oct; 29(46):8279-84.

Grimaldi-Bensouda L, Guillemot D, Godeau B, Benichou J, Lebrun-Frenay C, Papeix C, et al. [Autoimmune disorders and quadrivalent human papillomavirus vaccination of young female subjects](#). *J Intern Med*. 2014 Apr; 275(4):398-408. Epub 2013 Nov 22.

Klein NP, Hansen J, Chao C, Velicer C, Emery M, Slezak J, et al. [Safety of quadrivalent human papillomavirus vaccine administered routinely to females](#). *Arch Pediatr Adolesc Med*. 2012 Dec; 166(12):1140-8.

Macartney KK, Chiu C, Georgousakis M, Brotherton JM. [Safety of human papillomavirus vaccines: A review](#). *Drug Saf*. 2013 Jun; 36(6):393-412.

Markowitz LE, Dunne EF, Saraiya M, Chesson HW, Curtis CR, Gee J, et al. [Human papillomavirus vaccination: Recommendations of the Advisory Committee on Immunization Practices \(ACIP\)](#). *MMWR* 63(RR05);1-30

Moro PL, Zheteyeva Y, Lewis P, Shi J, Yue X, Museru OI, et al. [Safety of quadrivalent human papillomavirus vaccine \(Gardasil®\) in pregnancy: Adverse events among non-manufacturer reports in the Vaccine Adverse Event Reporting System, 2006-2013](#). *Vaccine*. 2015 Jan; 33(4): 519-22.

Scheller NM, Pasternak B, Svanström H, Hviid A. [Quadrivalent human papillomavirus vaccine and the risk of venous thromboembolism](#). *JAMA*. 2014 Jul;312(2):187-8.

Scheller NM, Svanström H, Pasternak B, Arnheim-Dahlström L, Sundström K, Katharina Fink K, et al. [Quadrivalent HPV vaccination and risk of multiple sclerosis and other demyelinating diseases of the central nervous system](#). *JAMA*. 2015 Jan; 313(1):54-61.

Slade BA, Leidel L, Vellozzi C, Woo EJ, Hua W, Sutherland A, et al. [Postlicensure safety surveillance for quadrivalent human papillomavirus recombinant vaccine](#). *JAMA*. 2009; 302(7):750-7

The Centers for Disease Control and Prevention, American Academy of Family Physicians, and American Academy of Pediatrics strongly recommend children receive all vaccines according to the recommended schedule.

Talk to your child's doctor or nurse about the vaccines recommended for their age.

	Flu <i>Influenza</i>	Tdap Tetanus, diphtheria, pertussis	HPV Human papillomavirus	Meningococcal		Pneumococcal	Hepatitis B	Hepatitis A		Inactivated Polio	MMR Measles, mumps, rubella	Chickenpox <i>Varicella</i>
				MenACWY	MenB							
7-8 Years	Green	Orange		Purple		Purple	Orange	Purple	Orange	Orange	Orange	Orange
9-10 Years	Green	Orange	Purple, Blue	Purple		Purple	Orange	Purple	Orange	Orange	Orange	Orange
11-12 Years	Green	Green, Orange	Green	Green, Orange		Purple	Orange	Purple	Orange	Orange	Orange	Orange
13-15 Years	Green	Orange	Orange	Orange		Purple	Orange	Purple	Orange	Orange	Orange	Orange
16-18 Years	Green	Orange	Orange	Green, Orange		Purple	Orange	Purple	Orange	Orange	Orange	Orange

More information:

Preteens and teens should get a flu vaccine every year.

Preteens and teens should get one shot of Tdap at age 11 or 12 years.

All 11-12 year olds should get a 2-shot series of HPV vaccine at least 6 months apart. A 3-shot series is needed for those with weakened immune systems and those age 15 or older.

All 11-12 year olds should get a single shot of a quadrivalent meningococcal conjugate vaccine (MenACWY). A booster shot is recommended at age 16.

Teens, 16-18 years old, **may** be vaccinated with a MenB vaccine.

 These shaded boxes indicate when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.

 These shaded boxes indicate the vaccine should be given if a child is catching-up on missed vaccines.

 These shaded boxes indicate the vaccine is recommended for children with certain health or lifestyle conditions that put them at an increased risk for serious diseases. See vaccine-specific recommendations at www.cdc.gov/vaccines/pubs/ACIP-list.htm.

 This shaded box indicates the vaccine is recommended for children not at increased risk but who wish to get the vaccine after speaking to a provider.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™



AMERICAN ACADEMY OF FAMILY PHYSICIANS

STRONG MEDICINE FOR AMERICA

Vaccine-Preventable Diseases and the Vaccines that Prevent Them

Diphtheria (Can be prevented by Tdap vaccination)

Diphtheria is a very contagious bacterial disease that affects the respiratory system, including the lungs. Diphtheria bacteria can be passed from person to person by direct contact with droplets from an infected person's cough or sneeze. When people are infected, the bacteria can produce a toxin (poison) in the body that can cause a thick coating in the back of the nose or throat that makes it hard to breathe or swallow. Effects from this toxin can also lead to swelling of the heart muscle and, in some cases, heart failure. In serious cases, the illness can cause coma, paralysis, and even death.

Hepatitis A (Can be prevented by HepA vaccination)

Hepatitis A is an infection in the liver caused by hepatitis A virus. The virus is spread primarily person-to-person through the fecal-oral route. In other words, the virus is taken in by mouth from contact with objects, food, or drinks contaminated by the feces (stool) of an infected person. Symptoms can include fever, tiredness, poor appetite, vomiting, stomach pain, and sometimes jaundice (when skin and eyes turn yellow). An infected person may have no symptoms, may have mild illness for a week or two, may have severe illness for several months, or may rarely develop liver failure and die from the infection. In the U.S., about 100 people a year die from hepatitis A.

Hepatitis B (Can be prevented by HepB vaccination)

Hepatitis B causes a flu-like illness with loss of appetite, nausea, vomiting, rashes, joint pain, and jaundice. Symptoms of acute hepatitis B include fever, fatigue, loss of appetite, nausea, vomiting, pain in joints and stomach, dark urine, grey-colored stools, and jaundice (when skin and eyes turn yellow).

Human Papillomavirus (Can be prevented by HPV vaccination)

Human papillomavirus is a common virus. HPV is most common in people in their teens and early 20s. It is the major cause of cervical cancer in women and genital warts in women and men. The strains of HPV that cause cervical cancer and genital warts are spread during sex.

Influenza (Can be prevented by annual flu vaccination)

Influenza is a highly contagious viral infection of the nose, throat, and lungs. The virus spreads easily through droplets when an infected person coughs or sneezes and can cause mild to severe illness. Typical symptoms include a sudden high fever, chills, a dry cough, headache, runny nose, sore throat, and muscle and joint pain. Extreme fatigue can last from several days to weeks. Influenza may lead to hospitalization or even death, even among previously healthy children.

Measles (Can be prevented by MMR vaccination)

Measles is one of the most contagious viral diseases. Measles virus is spread by direct contact with the airborne respiratory droplets of an infected person. Measles is so contagious that just

being in the same room after a person who has measles has already left can result in infection. Symptoms usually include a rash, fever, cough, and red, watery eyes. Fever can persist, rash can last for up to a week, and coughing can last about 10 days. Measles can also cause pneumonia, seizures, brain damage, or death.

Meningococcal Disease (Can be prevented by meningococcal vaccination)

Meningococcal disease is caused by bacteria and is a leading cause of bacterial meningitis (infection around the brain and spinal cord) in children. The bacteria are spread through the exchange of nose and throat droplets, such as when coughing, sneezing or kissing. Symptoms include sudden onset of fever, headache, and stiff neck. Meningococcal bacteria also cause blood infections. About one of every ten people who get the disease dies from it. Survivors of meningococcal disease may lose their arms or legs, become deaf, have problems with their nervous systems, become developmentally disabled, or suffer seizures or strokes.

Mumps (Can be prevented by MMR vaccination)

Mumps is an infectious disease caused by the mumps virus, which is spread in the air by a cough or sneeze from an infected person. A child can also get infected with mumps by coming in contact with a contaminated object, like a toy. The mumps virus causes swollen salivary glands under the ears or jaw, fever, muscle aches, tiredness, abdominal pain, and loss of appetite. Severe complications for children who get mumps are uncommon, but can include meningitis (infection of the covering of the brain and spinal cord), encephalitis (inflammation of the brain), permanent hearing loss, or swelling of the testes, which rarely results in decreased fertility.

Pertussis (Whooping Cough) (Can be prevented by Tdap vaccination)

Pertussis is caused by bacteria spread through direct contact with respiratory droplets when an infected person coughs or sneezes. In the beginning, symptoms of pertussis are similar to the common cold, including runny nose, sneezing, and cough. After 1-2 weeks, pertussis can cause spells of violent coughing and choking, making it hard to breathe, drink, or eat. This cough can last for weeks. Pertussis is most serious for babies, who can get pneumonia, have seizures, become brain damaged, or even die. About half of children under 1 year of age who get pertussis must be hospitalized.

Pneumococcal Disease (Can be prevented by pneumococcal vaccination)

Pneumonia is an infection of the lungs that can be caused by the bacteria called pneumococcus. This bacteria can cause other types of infections too, such as ear infections, sinus infections, meningitis (infection of the covering around the brain and spinal cord), and bacteremia (bloodstream infection). Sinus and ear infections are usually mild and are much more common than the

more serious forms of pneumococcal disease. However, in some cases pneumococcal disease can be fatal or result in long-term problems, like brain damage and hearing loss. Pneumococcal disease spreads when people cough or sneeze. Many people have the bacteria in their nose or throat at one time or another without being ill—this is known as being a carrier.

Polio (Can be prevented by IPV vaccination)

Polio is caused by a virus that lives in an infected person's throat and intestines. It spreads through contact with the stool of an infected person and through droplets from a sneeze or cough. Symptoms typically include sore throat, fever, tiredness, nausea, headache, or stomach pain. In about 1% of cases, polio can cause paralysis. Among those who are paralyzed, About 2 to 10 children out of 100 die because the virus affects the muscles that help them breathe.

Rubella (German Measles) (Can be prevented by MMR vaccination)

Rubella is caused by a virus that is spread through coughing and sneezing. In children rubella usually causes a mild illness with fever, swollen glands, and a rash that lasts about 3 days. Rubella rarely causes serious illness or complications in children, but can be very serious to a baby in the womb. If a pregnant woman is infected, the result to the baby can be devastating, including miscarriage, serious heart defects, mental retardation and loss of hearing and eye sight.

Tetanus (Lockjaw) (Can be prevented by Tdap vaccination)

Tetanus is caused by bacteria found in soil, dust, and manure. The bacteria enters the body through a puncture, cut, or sore on the skin. When people are infected, the bacteria produce a toxin (poison) that causes muscles to become tight, which is very painful. Tetanus mainly affects the neck and belly. This can lead to "locking" of the jaw so a person cannot open his or her mouth, swallow, or breathe. Complete recovery from tetanus can take months. One to two out of 10 people who get tetanus die from the disease.

Varicella (Chickenpox) (Can be prevented by varicella vaccination)

Chickenpox is caused by the varicella zoster virus. Chickenpox is very contagious and spreads very easily from infected people. The virus can spread from either a cough, sneeze. It can also spread from the blisters on the skin, either by touching them or by breathing in these viral particles. Typical symptoms of chickenpox include an itchy rash with blisters, tiredness, headache and fever. Chickenpox is usually mild, but it can lead to severe skin infections, pneumonia, encephalitis (brain swelling), or even death.

If you have any questions about your child's vaccines, talk to your healthcare provider.