

# COVID-19: Impact of Childhood Obesity on Health Outcomes

A Resource for Improving Measurable Impact

February 2, 2022

## Key Findings:

### Childhood obesity is linked to worse outcomes from COVID-19

- Obesity is a risk factor for hospitalization and mechanical ventilation from COVID-19 among children.<sup>1-3</sup> One study demonstrated that obesity was the **strongest risk factor for mechanical ventilation** among children ages two years or older with COVID-19.<sup>3</sup>

### Prevalence of childhood obesity is growing across the U.S.<sup>4</sup>

- Texas has the **8<sup>th</sup> highest rate** of childhood and adolescent (ages 10-17) obesity according to the most recent (2019-2020) national data estimate from the National Survey of Children's Health.<sup>5,6</sup>

### Minority youth are at risk

- In Texas, Hispanic and Black children have nearly twice the rate of obesity compared to non-Hispanic white children.<sup>7</sup>

### Stay-at-home orders impact healthy habits

- During stay-at-home orders, youth reported more physical inactivity, sedentary behavior, and consumption of unhealthy foods and sugar-sweetened beverages – all of which increase risk for weight gain.<sup>8-10</sup>

### Health behaviors persist into adulthood

- Poor health behaviors in childhood increase lifetime risk for serious conditions like obesity and diabetes.<sup>10,11</sup>

### Telemedicine can reduce barriers to care during the pandemic

- Chronic disease management requires frequent interactions with healthcare teams. Telemedicine ensures continuity of care for youth with obesity.<sup>12</sup>

## Impact of Obesity on COVID-19 Outcomes:

- As of January 27, 2022, children and adolescents accounted for 18.6% (11,411,047/61,294,2910) of U.S. COVID-19 cases.<sup>13</sup>
- As of November 20, 2021 the Centers for Disease Control and Prevention (CDC) reported that among 5,344 children hospitalized with COVID-19 and known condition status, 54.3% had an underlying health condition. The most common conditions included obesity (35.3%), asthma (15.8%), and neurologic conditions (14.3%)<sup>14</sup>
- Severe obesity is an even higher risk for COVID-19 complications among children.<sup>15</sup>

## Impact of COVID-19 on Obesity:

Studies indicate the pandemic and stay-at-home orders are impacting child health behaviors related to obesity.<sup>9</sup>

- During stay-at-home orders, children and adolescents:<sup>8,10</sup>
  - Decreased time spent playing sports and participating in physical activity
  - Increased sedentary time and screen time
  - Increased consumption of sugar-sweetened beverages and unhealthy foods
- Socioeconomic inequities may be exacerbated by COVID-19, as many families have limited resources available to purchase healthy foods and exercise at home during lockdowns.<sup>10,16,17</sup>
- A recent study conducted by the CDC found that the rate of increase in body mass index (BMI) among children and adolescents ages 2-19 years doubled during the pandemic (March-November 2020) compared to before the pandemic (January 2018-February 2020).<sup>18</sup>

## Mitigation & Prevention Recommendations:

### **Improve access to healthcare:**

- Enhance telehealth infrastructure for screening, prevention and treatment of obesity.<sup>19</sup> Telemedicine can be an effective approach for multidisciplinary teams to provide care to youth with obesity – especially during COVID-19 – by reducing travel barriers, building trust with healthcare providers, and ensuring continuity of care.<sup>9,12</sup>

### **Promote healthy habits:**

- Strengthen nutrition education, opportunities for physical activity, and availability of healthy foods in schools, early childhood, and afterschool programs.<sup>5,6,19,20</sup>
- Expand availability of resources to support the use of evidence-based childhood obesity prevention programs.<sup>6,19,20</sup>
- At the individual level, children and adolescents should participate in 60 minutes of physical activity each day, eat a variety of healthy foods, limit sedentary screen time, and get enough sleep each night to promote physical and emotional health.<sup>6,9,21-24</sup>

## Summary:

Though COVID-19 infections remain low in younger populations, children with obesity are at greater risk for hospitalization and mechanical ventilation from COVID-19. Moreover, stay-at-home orders are affecting health behaviors associated with weight gain and obesity among children, including poor dietary habits and physical inactivity. Recommendations include supporting telemedicine to coordinate care for youth with obesity, strengthening school- and community-based efforts to promote healthy eating and physical activity, and reinforcing COVID-19 prevention efforts in childcare centers and schools.

### TX RPC Network Member Content Experts

**Deanna Hoelscher, PhD, RDN, LD, CNS, FISBNPA**

The University of Texas Health Science Center at Houston (UTHealth) School of Public Health in Austin

**Alexandra van den Berg, PhD, MPH**

The University of Texas Health Science Center at Houston (UTHealth) School of Public Health in Austin

**Shreela Sharma, PhD, RD, LD**

The University of Texas Health Science Center at Houston (UTHealth) School of Public Health in Houston

**Sarah Messiah, PhD, MPH**

The University of Texas Health Science Center at Houston (UTHealth) School of Public Health in Dallas

**Emily Dhurandhar, PhD**

Metabologix, LLC

**Belinda Reininger, DrPH**

The University of Texas Health Science Center at Houston (UTHealth) School of Public Health in Brownsville

**Leah Whigham, PhD, FTOS**

The University of Texas Health Science Center at Houston (UTHealth) School of Public

## References:

1. Derespina KR, Kaushik S, Plichta A, et al. Clinical manifestations and outcomes of critically ill children and adolescents with coronavirus disease 2019 in new york city. *J Pediatr*. 2020. PMID: 32681989
2. Kim L, Whitaker M, O'Halloran A, et al. Hospitalization Rates and Characteristics of Children Aged <18 Years Hospitalized with Laboratory-Confirmed COVID-19 - COVID-NET, 14 States, March 1-July 25, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(32):1081-1088. Published 2020 Aug 14. PMID: 32790664
3. Zachariah P, Johnson CL, Halabi KC, et al. Epidemiology, Clinical Features, and Disease Severity in Patients With Coronavirus Disease 2019 (COVID-19) in a Children's Hospital in New York City, New York [published correction appears in *JAMA Pediatr*. 2021 Jun 21;:null]. *JAMA Pediatr*. 2020;174(10):e202430. PMID: 32492092
4. An R. Projecting the impact of the coronavirus disease-2019 pandemic on childhood obesity in the United States: A microsimulation model. *J Sport Health Sci*. 2020;9(4):302-312. PMID: 32454174
5. The Robert Wood Johnson Foundation and Trust for America's Health. (2021). The State of Obesity in Texas. State of Childhood Obesity website. <https://stateofchildhoodobesity.org/states/tx/>.
6. Trust for America's Health. The State of Obesity 2020: Better Policies for a Healthier America. Trust for America's Health website. <https://www.tfah.org/report-details/state-of-obesity-2020/>
7. The Michael & Susan Dell Center for Healthy Living. Healthy Children, Healthy State: Child Obesity Crisis in Texas. UTHealth School of Public Health website. <https://sph.uth.edu/research/centers/dell/resources/Child%20Obesity%20Crisis%20FINAL.pdf>.
8. Pietrobelli A, Pecoraro L, Ferruzzi A, et al. Effects of COVID-19 Lockdown on Lifestyle Behaviors in Children with Obesity Living in Verona, Italy: A Longitudinal Study. *Obesity (Silver Spring)*. 2020;28(8):1382-1385. PMID: 32352652
9. Silva CA, Queiroz LB, Fonseca CB, Silva LEVD, Lourenço B, Marques HHS. Spotlight for healthy adolescents and adolescents with preexisting chronic diseases during the COVID-19 pandemic. *Clinics (Sao Paulo)*. 2020;75:e1931. PMID: 32401965
10. Dunton GF, Do B, Wang SD. Early effects of the COVID-19 pandemic on physical activity and sedentary behavior in children living in the U.S. *BMC Public Health*. 2020;20(1):1351. Published 2020 Sep 4. PMID: 32887592
11. Rundle AG, Park Y, Herbstman JB, Kinsey EW, Wang YC. COVID-19-Related School Closings and Risk of Weight Gain Among Children. *Obesity (Silver Spring)*. 2020;28(6):1008-1009. PMID: 3222767

12. O'Hara VM, Johnston SV, Browne NT. The paediatric weight management office visit via telemedicine: pre- to post-COVID-19 pandemic. *Pediatr Obes*. July 6, 2020; PMID: [32627434](#)
13. American Academy of Pediatrics and the Children's Hospital Association. Children and COVID-19: State Data Report. American Academy of Pediatrics website. February 1, 2022. <https://www.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-covid-19-state-level-data-report/>.
14. Centers for Disease Control and Prevention (CDC). COVID-NET: A Weekly Summary of U.S. COVID-19 Hospitalization Data. Laboratory-Confirmed COVID-19-Associated Hospitalizations. Centers for Disease Control and Prevention website. February 2, 2022. [https://gis.cdc.gov/grasp/COVIDNet/COVID19\\_5.html](https://gis.cdc.gov/grasp/COVIDNet/COVID19_5.html).
15. Messiah SE, Xie L, Atem F, et al. Disparity Between United States Adolescent Class II and III Obesity Trends and Bariatric Surgery Utilization, 2015-2018 [published online ahead of print, 2020 Sep 15]. *Ann Surg*. PMID: 32941272
16. Sallis JF, Adlaka D, Oyeyemi A, Salvo D. An international physical activity and public health research agenda to inform coronavirus disease-2019 policies and practices. *J Sport Health Sci*. 2020;9(4):328-334. PMID: 32450160
17. Van Lancker W, Parolin Z. COVID-19, school closures, and child poverty: a social crisis in the making. *Lancet Public Health*. 2020;5(5):e243-e244. PMID: 32275858
18. Lange SJ, Kompaniyets L, Freedman DS, Kraus EM, Porter R, Blanck HM, Goodman AB. Longitudinal Trends in Body Mass Index Before and During the COVID-19 Pandemic Among Persons Aged 2-19 Years - United States, 2018-2020. *MMWR Morb Mortal Wkly Rep* 2021;70:1278-1283. DOI: <http://dx.doi.org/10.15585/mmwr.mm7037a3>.
19. Texas Department of State Health Services. Department of State Health Services Obesity Prevention Program Priority Objectives 2016-2021. Texas Health and Human Services website. 2018. [https://www.dshs.texas.gov/obesity/pdf/OPP\\_StratPlan\\_01032018.pdf](https://www.dshs.texas.gov/obesity/pdf/OPP_StratPlan_01032018.pdf)
20. Staples, T., Lakey, D., & Williams, M. A Report to the Texas Legislature from the Interagency Obesity Council. 2015. [https://dshs.texas.gov/uploadedFiles/Content/Prevention\\_and\\_Preparedness/obesity/2015\\_InteragencyObesity\\_Report\\_Final.pdf](https://dshs.texas.gov/uploadedFiles/Content/Prevention_and_Preparedness/obesity/2015_InteragencyObesity_Report_Final.pdf)
21. Calcaterra V, Vandoni M, Pellino VC, Cena H. Special Attention to Diet and Physical Activity in Children and Adolescents With Obesity During the Coronavirus Disease-2019 Pandemic. *Front Pediatr*. 2020;8:407. Published 2020 Jun 26. PMID: 32676491
22. World Health Organization. #Healthy at Home – Physical Activity. World Health Organization website. <https://www.who.int/news-room/campaigns/connecting-the-world-to-combat-coronavirus/healthyathome/healthyathome---physical-activity>
23. Nagata JM, Abdel Magid HS, Pettee Gabriel K. Screen Time for Children and Adolescents During the Coronavirus Disease 2019 Pandemic. *Obesity (Silver Spring)*. 2020;28(9): PMID: 32463530
24. Ribeiro KDDS, Garcia LRS, Dametto JFDS, Assunção DGF, Maciel BLL. COVID-19 and Nutrition: The Need for Initiatives to Promote Healthy Eating and Prevent Obesity in Childhood. *Child Obes*. 2020;16(4):235-237. PMID: 32429705